

Signal Analysis & Communication ECE 355

Ch. 2.4: Linear Constant Coefficient Differential/ Difference Equation

Lecture 12

04-10-2023




Ch. 2.4. LINEAR CONSTANT COEFFICIENT DIFFERENTIAL / DIFFERENCE EQUATION (LCCDE)

- An important class of CT/DT systems is that for which the input & output are related through LCCDE.
- LCCDE describes wide variety of physical phenomenon.
 - Electrical circuit RLC.
 - mechanical system - mass & spring with applied force] CT
 - Accumulation of saving in bank account] DT
- We can solve LCCDE to find EXPLICIT expressions between input & output.
 - NOTE: later we will see another approach, very easy & the preferred way to compute outputs for these systems.

Example

- Consider an RL circuit:



- Can be described by LCCDE

$$L \frac{dy(t)}{dt} + R y(t) = x(t)$$

$$\frac{dy(t)}{dt} + 2y(t) = x(t) \quad - (1)$$

- Let's solve this LCCDE for finding $y(t)$ with given conditions:

$$x(t) = e^{3t} u(t) \quad \& \quad y(t) = 0 \text{ for } t < 0$$

$$y(t) = ? \text{ for } t > 0$$

- From the calculus course, we know that

$$y(t) = y_h(t) + y_p(t) \quad - (2)$$

- $y_h(t)$: homogeneous solution, a solution of:

$$\frac{dy_h(t)}{dt} + 2y_h(t) = 0 \quad - (3) \quad \text{"Natural Response"}$$

- $y_p(t)$: particular solution, a special solution due to the I/P.
"Forced Response".

- To find $y_h(t)$, guess the solution of the form:

$$y_h(t) = Ae^{st}, \quad \text{for all 't'}$$

* Guessed exp. as its diff./int. is of exp. form as well.

- Inserting in eqn (3)

$$As e^{st} + 2A e^{st} = 0$$

$$A e^{st} (s+2) = 0$$

$$s = -2$$

$$\therefore y_h(t) = A e^{-2t} \quad (\text{for any } A) \quad - (4)$$

- Now to find $y_p(t)$, guess

$$y_p(t) = B e^{3t} \quad \text{for } t > 0 \quad - (5)$$

* a sign. of the same form as the given I/P.

- Inserting in eqn (1) for finding $y_p(t)$

$$3B e^{3t} + 2B e^{3t} = e^{3t}$$

$$3B + 2B = 1$$

$$B = 1/5$$

$$\therefore \text{eqn. (5)} \Rightarrow y_{f(t)} = \frac{1}{5} e^{3t} \text{ for } t > 0 - (6)$$

- Substituting eqn. (4) & (6) in eqn (2)

$$y_{(t)} = A e^{-2t} + \frac{1}{5} e^{3t} - (7)$$

- To determine (A), we use the initial condition $y_{(0)} = 0$ for $t < 0$

$$\text{Eqn. (7)} \Rightarrow 0 = A + \frac{1}{5}$$

$$A = -\frac{1}{5}$$

$$\therefore \text{Therefore, eqn. (7)} \Rightarrow y_{(t)} = -\frac{1}{5} e^{-2t} + \frac{1}{5} e^{3t} \quad t > 0$$

NOTE

Suppose the initial condition was $y_{(0)} = 1$, then,

$$\text{eqn (7)} \Rightarrow 1 = A + \frac{1}{5}$$

$$A = \frac{4}{5}$$

$$\therefore \text{eqn (7)} \Rightarrow y_{(t)} = \frac{4}{5} e^{-2t} + \frac{1}{5} e^{3t}, \quad t > 0 - (8)$$

But in this case the initial rest condition is not satisfied for the considered LTI systems, i.e.,

$$\begin{array}{l|l} y_{(t)} = 0 & \text{for } t < t_0 \\ \text{if } x_{(t)} = 0 & \text{for } t < t_0 \end{array} \quad \Bigg| \quad \forall t, t_0$$

provides close approximation to most LTI systems!

For the (Causal) LTI systems, we consider in this course, the initial rest condition should be satisfied!

- The system defined by eqn (8) is either not causal or not an LTI.

* Remember (ref. - Lecture 10):

- Causality also ensures time invariance of LTI systems.
- That is, with zero initial values, the response at different times will be the same (with the same I/p) [assuming R & C do not change over time].

General Form of LCDE (CT)

$$\sum_{k=0}^N a_k \frac{d^k}{dt^k} y(t) = \sum_{k=0}^M b_k \frac{d^k}{dt^k} x(t) \quad - (A)$$

Standard Solution:

$$y(t) = y_h(t) + y_p(t)$$

Use roots of Charac. Eqn.

Guess!

$x(t)$	$y_p(t)$
e^{st}	Be^{st}
$\cos(\omega t)$	$A \cos(\omega t) + B \sin(\omega t)$
$\vdots$	$\vdots$

Auxiliary conditions: (Initial Conditions - Rest)

$$y(t_0) = \frac{d}{dt} y(t_0) = \dots = \frac{d^{N-1}}{dt^{N-1}} y(t_0) = 0$$

(assume continuity of $y(t)$ & its derivatives)

NOTE : We will see later that solving LCCDE is more straight forward in freq. domain.

LCCDE (DT)

- Recall that the differential in CT is replaced with the difference in DT.

- This implies that,

$$\frac{d}{dt} y(t) \rightarrow d^1 y[n] \triangleq y[n] - y[n-1]$$

$$\begin{aligned} \frac{d^2}{dt^2} y(t) &\rightarrow d^2 y[n] \triangleq dy[n] - dy[n-1] \\ &= (y[n] - y[n-1]) - (y[n-1] - y[n-2]) \\ &= y[n] - 2y[n-1] + y[n-2] \quad - \textcircled{B} \end{aligned}$$

$\vdots$

$$d^k y[n] = d^{k-1} y[n] - d^{k-1} y[n-1]$$

- Therefore, the DT analogous of eqn (A) can be written as:

$$\hat{a}_0 y[n] + \sum_{k=1}^N \hat{a}_k d^k y[n] = \hat{b}_0 x[n] + \sum_{k=1}^M \hat{b}_k d^k x[n] \quad - \textcircled{C}$$

- Equivalently (using eqn (B), eqn (C) can be expressed as:

$$\sum_{k=0}^N a_k y[n-k] = \sum_{k=0}^M b_k x[n-k] \quad - \textcircled{D}$$

Standard Solution

I. Similar to the CT case

$$y[n] = y_h[n] + y_p[n]$$

- assume initial rest conditions for causal LTI systems, that is,

$$\left. \begin{array}{l} y[n] = 0 \quad \text{for } n < n_0 \\ \text{if } x[n] = 0 \quad \text{for } n < n_0 \end{array} \right\} \forall n, n_0$$

II. We have eqn ① as

$$a_0 y[n] + \sum_{k=1}^N a_k y[n-k] = \sum_{k=0}^M b_k x[n-k]$$

$$y[n] = \frac{1}{a_0} \left[\sum_{k=0}^M b_k x[n-k] - \sum_{k=1}^N a_k y[n-k] \right]$$

- And then compute recursively!

Example

2.15 (textbook)

$$y[n] - \frac{1}{2} y[n-1] = x[n], \quad x[n] = k \delta[n]$$

$$y[n] = x[n] + \frac{1}{2} y[n-1]$$

$$y[0] = x[0] + \frac{1}{2} y[-1] = k + 0 = k$$

$$y[1] = x[1] + \frac{1}{2} y[0] = 0 + \frac{k}{2} = \frac{k}{2}$$

$\vdots$

$\vdots$

$\vdots$

$$y[n] = \dots$$

Causal LTI

$x[n] = 0$, for $n < -1$
 $\therefore y[n] = 0$, for $n < -1$
 $\Rightarrow y[-1] = 0$

→ Check yourselves!